

Applied Biological Materials Inc.

Tel: 1-866-757-2414
Email: info@abmGood.com
Website: www.abmGood.com

SafeView™ Nucleic Acid Stains

Store at 4°C

Cat. No.	Description	Quantity
G108	SafeView™ Classic	1.0 ml
G108-G	Safe-Green™	1.0 ml
G108-R	Safe-Red™	1.0 ml
G108-W	Safe-White™	1.0 ml
G108-P	Safe-Pack™	G108-G,R,W

Product Description

SafeViewTM products represent a new and safe class of nucleic acid stains for the visualization of double-stranded DNA, single-stranded DNA, and RNA in agarose and polyacrylamide gels. The dyes are developed to replace toxic Ethidium Bromide (EtBr, a potent mutagen), commonly used in gel electrophoresis for visualization of nucleic acids in agarose and polyacrylamide gels.

SafeViewTM products are non-carcinogenic by the Ames-test. The results are negative in both the mouse marrow chromophilous erythrocyte micronucleus and mouse spermary spermatocyte chromosomal aberration tests.

NOTE: SafeViewTM Nucleic Acid Stains are non-carcinogenic, but may cause skin and eye irritaions. Always wear gloves when working with the product.

$SafeView^{TM}\ Classic$ - Substitute for Ethidium Bromide in agarose gel

SafeViewTM Classic is used the same way as Ethidium Bromide in agarose and polyacrylamide gel electrophoresis. It emits green fluorescence when bound to dsDNA and ssDNA and red fluorescence when bound to RNA. This stain has two fluorescence excitation maxima when bound to nucleic acid, at approximately 290 nm and 490 nm.

Protocol

- 1. Prepare a 100 ml agarose or polyacrylamide solution.
- Add 5 µl SafeView™ Classic to the gel solution.
- 3. Mix gently; the solution should have no air bubbles.
- 4. For agarose gel, let the solution cool down to 60 70°C and cast the gel. Contd...

- For polyacrylamide gel, add APS and TEMED and cast the gel according to regular polyacrylamide gel casting protocol.
- 5. Run gel electrophoresis with 5 µl SafeView™ Classic per 100 ml buffer.
- 6. View the results under UV or blue LED light.

Safe-Green, Red, White, Pack™ - Substitute for loading dye

With SafeViewTM dyes (Safe-Green,Red,White,PackTM), you do not need to add any dyes to both gel matrix and running buffers. SafeViewTM dyes are provided in a form of 6X sample loading dyes and they are to be added to your samples only. The SafeViewTM dyes eliminate contamination risk of glassware or gel running tank as associated with EtBr. After the electrophoresis, view and document your results as you would do with EtBr staining protocols.

Protocol

- 1. Prepare a 100 ml agarose or polyacrylamide solution.
- 2. Mix gently without introducing any air bubbles.
- 3. For agarose gel, let the solution cool down to 60 70°C and cast the gel. For polyacrylamide gel, add APS and TEMED and cast the gel according to regular polyacrylamide gel casting protocol.
- 4. Mix samples and DNA marker with SafeViewTM dye at a1:5 (dye: sample) dilution rate.
- 5. Following electrophoresis, view the results under UV or blue LED light.