



## CheKine™ Tannase (TAN) Activity Colorimetric Assay Kit

Cat #: KTB1542

Size: 48 T/96 T

	<b>Tannase (TAN) Activity Colorimetric Assay Kit</b>		
<b>REF</b>	<b>Cat #:</b> KTB1542	<b>LOT</b>	<b>Lot #:</b> Refer to product label
	<b>Detection range:</b> 0.3125-20 µmol/mL (The Detection range corresponds to the standard)		<b>Sensitivity:</b> 0.156 µmol/mL (The Sensitivity corresponds to the standard)
	<b>Applicable samples:</b> Plant Tissues, Fungi, Bacteria		
	<b>Storage:</b> Stored at 4°C for 6 months		

### Assay Principle

Tannase, full name is Tannin Acyl Hydrolase (Tannase, EC 3.1.1.20). Tannase exists in tannin-rich plants and also widely exists in microorganisms. Tannase hydrolyses the ester bonds and depside bonds in gallic acid tannins to release gallic acid and glucose. The enzyme can be produced by molds such as *Aspergillus Niger*, *Aspergillus oryzae*. It can be used to treat tannin and protein in beer to make it clear and transparent. It can also be used to remove the astringency of persimmon and other products. And it can also be used to make instant tea to prevent turbid fermented tea. CheKine™ Tannase (TAN) Activity Colorimetric Assay Kit provides a convenient tool for detection of Tannase activity. The principle is to use propyl gallate (PG) as the substrate for the enzymatic reaction of tanninase, which has a characteristic absorption peak at 270 nm. The Tannase activity measure the change of the absorbance at 270 nm before and after the reaction, and calculate the Tannase activity of the sample can be calculated by measuring the absorbance at 270 nm.

### Materials Supplied and Storage Conditions

Kit components	Size		Storage conditions
	48 T	96 T	
Extraction Buffer	100 mL	100 mL×2	4°C
Substrate	5 mL	10 mL	4°C, protected from light
Standard	1	2	4°C, protected from light

### Materials Required but Not Supplied

- Microplate Reader capable of measuring absorbance at 270 nm
- Water Bath, Ice Maker, Refrigerated Centrifuge
- 96-well UV microplate, Precision Pipettes, Disposable Pipette Tips
- Anhydrous Ethanol
- Homogenizer (for Tissue Samples)

## Reagent Preparation

**Extraction Buffer:** Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

**Substrate:** Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C, protected from light.

**Standard:** Add 1.178 mL of Anhydrous Ethanol to dissolve before use. The concentration is 20 µmol/mL. This solution can be stored at 4°C for one week or stored at -20°C for long time.

**Standard curve setting:** Dilute 20 µmol/mL standard with Extraction Buffer as shown in the table below.

	Volume of Standard	Volume of Extraction Buffer (µL)	The concentration of Standard (µmol/mL)
Std.1	200 µL of 20 µmol/mL	0	20
Std.2	100 µL of Std.1(20 µmol/mL)	100	10
Std.3	100 µL of Std.2 (10 µmol/mL)	100	5
Std.4	100 µL of Std.3 (5 µmol/mL)	100	2.5
Std.5	100 µL of Std.4 (2.5 µmol/mL)	100	1.25
Std.6	100 µL of Std.5 (1.25 µmol/mL)	100	0.625
Std.7	100 µL of Std.6 (0.625 µmol/mL)	100	0.313

## Sample Preparation

**Note:** Fresh samples are recommended, If not assayed immediately, samples can be stored at -80°C for one month. Processed samples must be assayed immediately. If the protein concentration of the sample is need to determined, it is recommended to use Abbkine Cat #: KTD3001 Protein Quantification Kit (BCA Assay) to measure the protein concentration of the sample.

1. Plant Tissues: Weigh 0.1 g tissue, add 1 mL Extraction Buffer and homogenize on ice. Centrifuge at 10,000 g for 10 min at 4°C. Use supernatant for assay, and place it on ice to be tested.
2. Fungi or Bacteria: Collect  $5 \times 10^6$  Fungi or Bacteria into the centrifuge tube, wash Fungi or Bacteria with cold PBS, discard the supernatant after centrifugation; add 1 mL Extraction Buffer to ultrasonically disrupt the Fungi or Bacteria in ice bath 5 min (power 20% or 200 W, ultrasonic 3 s, interval 7 s, repeat 30 times). Centrifuge at 10,000 g for 10 min at 4°C. Use supernatant for assay, and place it on ice to be tested.

## Assay Procedure

1. Preheat the microplate reader for more than 30 min, and adjust the wavelength to 270 nm.
2. Add 50 µL of sample into an EP tube as a control tube, bath in boiling water for 5 min, and cool to room temperature.
3. Add the following reagents respectively into each EP tube :

	Blank tube (µL)	Standard tube (µL)	Test tube (µL)	Control tube (µL)
Extraction Buffer	200	150	100	100
Stds.	0	50	0	0
sample	0	0	50	50 (deactivated)
Substrate	0	0	50	50

Mix well, incubate at 40°C water bath for 10 min, and then immediately take a boiling water bath for 5 min. After cooling, centrifuge at 10,000 g at room temperature (25°C) for 10 min, and take the supernatant.

Add the following reagents to a 96-well UV microplate:

supernatant	10	10	10	10
Extraction Buffer	190	190	190	190

4. Mix well, read the values at 270 nm. Recorded as  $A_{Blank}$ ,  $A_{Standard}$ ,  $A_{Test}$  and  $A_{Control}$ , respectively. Finally, calculate  $\Delta A_{Test} = A_{Control} - A_{Test}$ ,  $\Delta A_{Standard} = A_{Standard} - A_{Blank}$ . (Only one blank well needs to be detected, a control is required for each sample).

**Note: In order to guarantee the accuracy of experimental results, need to do a pre-experiment with 2-3 samples. If  $\Delta A_{Test}$  is greater than 1.0, the sample can be appropriately diluted with Extraction Buffer, the calculated result multiplied by the dilution factor. If  $\Delta A_{Test}$  is less than 0.001, increase the sample quantity appropriately.**

## Data Analysis

**Note: We provide you with calculation formulae, including the derivation process and final formula. The two are exactly equal. It is suggested that the concise calculation formula in bold is final formula.**

### 1. Drawing of standard curve

With the concentration of the standard Solution as the y-axis and the  $\Delta A_{Standard}$  as the x-axis, draw the standard curve. Substitute the  $\Delta A_{Test}$  into the equation to obtain the y value ( $\mu\text{mol/mL}$ ).

### 2. Calculate the activity of Tannase (TAN)

#### (1) By sample weight

Unit Definition: 1 nmol PG reduced per min in 1 g tissue reaction system at 40°C is defined as a unit of enzyme activity.

$$\text{TAN (U/g)} = y \times 1,000 \times F \times V_{\text{Reaction Total}} \div (W \times V_{\text{Sample}} \div V_{\text{Sample Total}}) \div T \times n = \mathbf{8,000 \times y \div W \times n}$$

#### (2) By cells number of Fungi or Bacteria

Unit Definition: 1 nmol PG reduced per min in  $10^4$  cells number of Fungi or Bacteria reaction system at 40°C is defined as a unit of enzyme activity.

$$\text{TAN (U/}10^4 \text{ cells)} = y \times 1,000 \times F \times V_{\text{Reaction Total}} \div (V_{\text{Sample}} \times 500 \div V_{\text{Sample Total}}) \div T \times n = \mathbf{16 \times y \times n}$$

#### (3) By protein concentration


Unit Definition: 1 nmol PG reduced per min in 1 mg tissue protein reaction system at 40°C is defined as a unit of enzyme activity.

$$\text{TAN (U/mg prot)} = y \times 1,000 \times F \times V_{\text{Reaction Total}} \div (C_{\text{pr}} \times V_{\text{Sample}}) \div T \times n = \mathbf{8,000 \times y \div C_{\text{pr}} \times n}$$

Where: 1,000: 1  $\mu\text{mol} = 1,000 \text{ nmol}$ ; F: Dilution factor of supernatant,  $F = 200 \mu\text{L} \div 10 \mu\text{L} = 20$ ;  $V_{\text{Reaction Total}}$ : total reaction volume, 0.2 mL; W: sample weight, g;  $V_{\text{sample}}$ : sample volume added, 0.05 mL;  $V_{\text{Sample Total}}$ : Extraction Buffer volume added, 1 mL; T: reaction time, 10 min; n: dilution multiple of sample; 500: Total cells number of Fungi or Bacteria,  $5 \times 10^6$ ; Cpr: sample protein concentration, mg/mL.

## Typical Data

Typical standard curve



3/4

Version 20220421

Figure 1. Standard Curve of PG in 96-well plate assay-data provided for demonstration purposes only. A new standard Curve must be generated for each assay.

## Recommended Products

Catalog No.	Product Name
KTB1540	CheKine™ Plant Total Phenols (TP) Colorimetric Assay Kit
KTB1530	CheKine™ Plant Flavonoids Colorimetric Assay Kit
KTB1541	CheKine™ Tannin Colorimetric Assay Kit

## Disclaimer

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.