


Thiazolyl Blue

Cat. No.:	HY-15924
CAS No.:	298-93-1
Molecular Formula:	C ₁₈ H ₁₆ BrN ₅ S
Molecular Weight:	414.32
Target:	Fluorescent Dye
Pathway:	Others
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro

DMSO : 25 mg/mL (60.34 mM; ultrasonic and warming and heat to 60°C)
H₂O : 0.88 mg/mL (2.12 mM; Need ultrasonic and warming)

Concentration	Solvent	Mass	1 mg			5 mg			10 mg		
			Concentration			Concentration			Concentration		
1 mM			2.4136 mL			12.0680 mL			24.1359 mL		
5 mM			0.4827 mL			2.4136 mL			4.8272 mL		
10 mM			0.2414 mL			1.2068 mL			2.4136 mL		

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: PBS
Solubility: 6.67 mg/mL (16.10 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (5.02 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.08 mg/mL (5.02 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Thiazolyl Blue (MTT) is a cell-permeable and positively charged tetrazolium dye that is used to detect reductive metabolism in cells. Thiazolyl Blue is taken up by cells through the plasma membrane and then reduced to formazan by intracellular NAD (P) H-oxidoreductases. Thiazolyl Blue is frequently used in colorimetric assays to measure cell proliferation, cytotoxicity, and apoptosis^[1]. Storage: protect from light.

In Vitro

- Preparation of MTT working solution
Dissolve MTT by PBS to obtain 5 mg/mL of MTT.

2. Cell proliferation detection (96-well plates)

2.1 Inoculated cells: Prepared single cell suspension to prepared with culture medium containing 10% FBS and inoculated into 96-well plate with 1000-10000 cells per well, volume of 100 μ L per well.

2.2 Incubated cells: 37 ^\circ C, 5% CO₂, Incubate for 24-72 h.

2.3 Add 10 μ L MTT in per well and incubate for 4 h, Discard the supernatant, For suspended cells, need centrifuge firstly.

2.4 Add 100 μ L DMSO, shock 10 min to dissolve the crystal completely

2.5 Monitor the absorbance increase with an absorbance plate reader at OD = 562 nm.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell Mol Immunol. 2021 Mar;18(3):613-620.
- Nat Commun. 2022 Nov 10;13(1):6796.
- Bioact Mater. 2020 Mar 30;5(2):398-409.
- Small. 2023 Jan 12;e2207194.
- J Hazard Mater. 2020 Apr 5;387:121686.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Berridge MV, et al. Tetrazolium dyes as tools in cell biology: new insights into their cellular reduction. Biotechnol Annu Rev. 2005;11:127-52.

[2]. Berridge MV, et al. Tetrazolium dyes as tools in cell biology: new insights into their cellular reduction. Biotechnol Annu Rev. 2005;11:127-52.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA