


## Cefotaxime sodium

|                    |                                                                                                                                |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------|
| Cat. No.:          | HY-A0088                                                                                                                       |
| CAS No.:           | 64485-93-4                                                                                                                     |
| Molecular Formula: | C <sub>16</sub> H <sub>16</sub> N <sub>5</sub> NaO <sub>7</sub> S <sub>2</sub>                                                 |
| Molecular Weight:  | 477.45                                                                                                                         |
| Target:            | Bacterial; Antibiotic; Beta-lactamase                                                                                          |
| Pathway:           | Anti-infection                                                                                                                 |
| Storage:           | 4°C, sealed storage, away from moisture<br>* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture) |



### SOLVENT & SOLUBILITY

|                                                                               |                                                                                                                                                   |                                                      |           |            |            |
|-------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------|-----------|------------|------------|
| In Vitro                                                                      | H <sub>2</sub> O : 50 mg/mL (104.72 mM; Need ultrasonic)<br>DMSO : 45 mg/mL (94.25 mM; Need ultrasonic)                                           |                                                      |           |            |            |
|                                                                               | Preparing Stock Solutions                                                                                                                         | <div>Solvent<br/>Concentration</div> <div>Mass</div> | 1 mg      | 5 mg       | 10 mg      |
|                                                                               |                                                                                                                                                   | 1 mM                                                 | 2.0945 mL | 10.4723 mL | 20.9446 mL |
|                                                                               |                                                                                                                                                   | 5 mM                                                 | 0.4189 mL | 2.0945 mL  | 4.1889 mL  |
|                                                                               |                                                                                                                                                   | 10 mM                                                | 0.2094 mL | 1.0472 mL  | 2.0945 mL  |
| Please refer to the solubility information to select the appropriate solvent. |                                                                                                                                                   |                                                      |           |            |            |
| In Vivo                                                                       | 1. Add each solvent one by one: PBS<br>Solubility: 100 mg/mL (209.45 mM); Clear solution; Need ultrasonic                                         |                                                      |           |            |            |
|                                                                               | 2. Add each solvent one by one: 10% DMSO >> 90% corn oil<br>Solubility: ≥ 2.5 mg/mL (5.24 mM); Clear solution                                     |                                                      |           |            |            |
|                                                                               | 3. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline<br>Solubility: ≥ 2.08 mg/mL (4.36 mM); Clear solution         |                                                      |           |            |            |
|                                                                               | 4. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)<br>Solubility: 2.08 mg/mL (4.36 mM); Suspended solution; Need ultrasonic |                                                      |           |            |            |

### BIOLOGICAL ACTIVITY

|                           |                                                                                                                                                                                                                                                        |          |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|
| Description               | Cefotaxime (Cefotaxim) sodium, a β-lactamase stable cephalosporin and a third-generation cephalosporin antibiotic, possesses broad-spectrum antibiotic activity against numerous Gram-positive and Gram-negative bacteria <sup>[1][2][3][4][5]</sup> . |          |
| IC <sub>50</sub> & Target | β-lactam                                                                                                                                                                                                                                               | β-lactam |
| In Vitro                  | Cefotaxime sodium exhibits an MIC of 0.0625 mg/L for <i>V. vulnificus</i> CMCP6 <sup>[4]</sup> .                                                                                                                                                       |          |

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

The combination of ciprofloxacin and cefotaxime is more effective in clearing *V. vulnificus* in vivo than previously used regimens<sup>[4]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

|                 |                                                                                                                                                                                       |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Animal Model:   | Female, specific pathogen free, 8-week-old BALB/c mice <sup>[4]</sup> .                                                                                                               |
| Dosage:         | 30 mg/kg.                                                                                                                                                                             |
| Administration: | IP every 6 h.                                                                                                                                                                         |
| Result:         | The viable bacterial counts in liver were lower in mice treated with cefotaxime-plus-ciprofloxacin than in those treated with cefotaxime alone ( $P < 0.001$ at 24 h and 48 h, each). |

## REFERENCES

- [1]. Gums JG, et al. Differences between ceftriaxone and cefotaxime: microbiological inconsistencies. *Ann Pharmacother*. 2008 Jan;42(1):71-9.
- [2]. Woodfield JC, et al. A comparison of the prophylactic efficacy of ceftriaxone and cefotaxime in abdominal surgery. *Am J Surg*. 2003 Jan;185(1):45-9.
- [3]. Scholz H, et al. Prospective comparison of ceftriaxone and cefotaxime for the short-term treatment of bacterial meningitis in children. *Chemotherapy*. 1998 Mar-Apr;44(2):142-7.
- [4]. E L Francke, et al. Use of cefotaxime, a beta-lactamase stable cephalosporin, in the therapy of serious infections, including those due to multiresistant organisms. *Am J Med*. 1981 Sep;71(3):435-42.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA