

Thrombin (MW 37kDa)

Cat. No.:	HY-114164
CAS No.:	9002-04-4
Molecular Weight:	37000(Average)
Target:	Thrombin
Pathway:	Metabolic Enzyme/Protease
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.

Thrombin (MW 37kDa)

BIOLOGICAL ACTIVITY

Description	Thrombin (MW 37kDa) is a Na ⁺ -activated, allosteric serine protease that plays opposing functional roles in blood coagulation. Thrombin recognition sequence and can be used to digest GST-tagged proteins.
In Vitro	p62 is cleaved from the agarose beads as well as the GST tag by addition of Thrombin (1 Unit/100 µg protein) at room temperature for 20 h^[3]. Thrombin solution: Dissolve in PBS prechilled to 4°C. Swirl gently. Store solution in small aliquots at -80°C to preserve activity. Cleavage buffer: PBS (140 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 1.8 mM KH₂PO₄) pH 7.3. Elution buffer: 50 mM Tris-HCl, 10 mM reduced glutathione, pH 8.0 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cancer Commun (Lond). 2025 Feb;45(2):143-166.
- Allergy. 2022 Jul;77(7):2104-2120.
- Front Pharmacol. 2022 Jan 10;12:792263.
- Antiviral Res. 2023 Jun:214:105606.
- Mol Carcinog. 2024 Jul;63(7):1288-1302.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Geetha T, et al. p62 serves as a shuttling factor for TrkA interaction with the proteasome. Biochem Biophys Res Commun. 2008 Sep 12;374(1):33-7.
- [2]. Golderman V, et al. Thrombin and the Protease-Activated Receptor-1 in Organophosphate-Induced Status Epilepticus. J Mol Neurosci. 2018 Dec 4.
- [3]. Krenzlin H, et al. The Importance of Thrombin in Cerebral Injury and Disease. Int J Mol Sci. 2016 Jan 11;17(1).

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite F, Monmouth Junction, NJ 08852, USA